



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $y^2 = m^2 x$ be the equation of the curve. In this case we have,

$$\bar{x} = \frac{\int_0^a \int_0^{mx^{\frac{1}{2}}} (a-x) x dx dy}{\int_0^a \int_0^{mx^{\frac{1}{2}}} (a-x) dx dy} = \frac{\int_0^a (a-x) x^{\frac{3}{2}} dx}{\int_0^a (a-x) x^{\frac{1}{2}} dx} = \left[\frac{\frac{2}{5} a x^{\frac{5}{2}} - \frac{2}{7} x^{\frac{7}{2}}}{\frac{2}{3} a x^{\frac{3}{2}} - \frac{2}{5} a x^{\frac{5}{2}}} \right]_0^a = \frac{3}{7} a.$$

Therefore, the center of pressure is $\frac{4}{7}a$ below the surface.

(2). To find the center of pressure on the outer figure. We have

$$\begin{aligned} \bar{x} &= \frac{\int_0^a \int_0^{b-mx^{\frac{1}{2}}} (a-x) x dx dy}{\int_0^a \int_0^{b-mx^{\frac{1}{2}}} (a-x) dx dy} = \frac{\int_0^a (a-x)(b-mx^{\frac{1}{2}}) x dx}{\int_0^a (a-x)(b-mx^{\frac{1}{2}}) dx} \\ &= \left[\frac{\frac{1}{2} a b x^2 - \frac{2}{5} a m x^{\frac{5}{2}} - \frac{1}{3} b x^3 + \frac{2}{7} m x^{\frac{7}{2}}}{a b x - \frac{2}{3} a m x^{\frac{3}{2}} - \frac{1}{2} b x^2 + \frac{2}{5} m x^{\frac{5}{2}}} \right]_0^a \end{aligned}$$

But the curve gives $b^2 = m^2 a$ or $m = b/a^{\frac{1}{2}}$. Substituting this value of m , and reducing, we have, $\bar{x} = \frac{11}{49}a$. Hence the depth of the center of pressure $= \frac{3}{4}\frac{8}{9}a$.

Also solved by G. B. M. ZERR, and T. T. DAVIS.

DIOPHANTINE ANALYSIS.

89. Proposed by J. H. DRUMMOND, LL. D., Portland, Me.

Show that in $2x^2 + 2y^2 - z^2 = \square \dots (1)$,

$2x^2 + 2z^2 - y^2 = \square \dots (2)$,

$2y^2 + 2z^2 - x^2 = \square \dots (3)$,

any two numbers and their sum and difference will satisfy the conditions.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let $2x^2 + 2y^2 - z^2 = a^2$, $2x^2 + 2z^2 - y^2 = b^2$, $2y^2 + 2z^2 - x^2 = c^2$.

Adding, $2x^2 + 2y^2 + 2z^2 = \frac{2}{3}(a^2 + b^2 + c^2)$.

$\therefore z = \pm \frac{1}{3} \sqrt{(2b^2 + 2c^2 - a^2)}$,

$y = \pm \frac{1}{3} \sqrt{(2a^2 + 2c^2 - b^2)}$,

$x = \pm \frac{1}{3} \sqrt{(2a^2 + 2b^2 - c^2)}$.

Let $b = (n+1)a$, $c = na$.

$\therefore z = \pm \frac{a}{3} (2n+1)$, $y = \pm \frac{a}{3} (n-1)$, $x = \pm \frac{a}{3} (n+2)$.

Solutions of problem 87 were received from H. S. VANDIVER, G. B. M. ZERR, J. H. DRUMMOND, and H. C. WHITAKER; of problem 88, from G. B. M. ZERR, J. H. DRUMMOND, J. SCHEFFER, and H. C. WHITAKER.